arXiv:1210.0931v1 [math.AP] 2 Oct 2012

THE RIEMANN-HILBERT PROBLEM FOR ELLIPTIC VECTOR FIELDS WITH DEGENERACIES

A. AINOUZ[1], K. E. BOUTARENE[1], A. MEZIANI[2]

Abstract. The boundary value problem

$$Lu = f \ \text{ in } \Omega\,, \quad \mathrm{Re}(\overline{\Lambda}\, u) = \phi \ \text{ on } \partial\Omega\,,$$

is studied for a class of planar complex vector fields L in a simply connected open set $\Omega \subset \mathbb{R}^2$. The first integrals of L are used to reduce the problem into a collection of classical Riemann-Hilbert problems with discontinuous data.

Introduction

This paper deals with boundary value problems for planar elliptic vector fields with degeneracies. The vector field L is nonsingular, complex valued, and real analytic in an open set containing $\overline{\Omega}$, where Ω is open, bounded, simply connected, and has a smooth boundary. We assume that L is locally solvable and is elliptic except on the analytic variety Σ given by $L \wedge \overline{L} = 0$. The boundary value problems considered here are versions of the Riemann-Hilbert problem (RH problem for short) for the vector field L:

$$\left\{ \begin{array}{lll} Lu = f & \text{in} & \Omega\,, \\ \mathrm{Re}(\overline{\Lambda}\, u) = \phi & \text{on} & \partial\Omega\,, \end{array} \right. \tag{0.1}$$

where $\Lambda\,,\ \phi$ are Hölder continuous on $\partial\Omega$ with $|\Lambda| = 1$, ϕ is $\mathbb{R}$-valued, and $f \in C^\infty(\overline{\Omega})$.

The analytic variety Σ consists of the set Σ^0, of points of finite type, and the set Σ^∞, of points of infinite type (see Section 1 for definitions). Σ^∞ is a one-dimensional submanifold in $\overline{\Omega}$ and L is tangent to Σ^∞. The connected components of Σ^∞ are the (one-dimensional) orbits of L. We assume that these orbits are minimal (to be defined below). Let $\Gamma_1, \cdots, \Gamma_N$ be the non closed orbits of L such that each Γ_j connects in Ω two distinct points of $\partial\Omega$. These orbits play a crucial role in the solvability of Problem (0.1). For a given function Λ, defined on $\partial\Omega$, we associate an index $\kappa_j \in \mathbb{Z}$ in each connected component Ω_j of $\Omega\backslash\Gamma_1 \cup \cdots \cup \Gamma_N$, and to each orbit Γ_j with ends $p_j^-\,,\, p_j^+$ on $\partial\Omega$, we attach a unique real number $\alpha_j \in [0,\ 1)$. This number measures, in some sense, the jump of Λ along the orbit.

In general, it is not possible to find continuous solutions of Problem (0.1) on $\overline{\Omega}$, even in simple cases. The reason is that some of the indices κ_j might be negative. We will then allow for solutions to have isolated singular points: at most one singular point in each component Ω_j with index $\kappa_j < 0$. In fact, the solution u is continuous on $\overline{\Omega}\backslash\{p_1, \cdots, p_n\}$ and, near each singular point p_j, it behaves (through first integrals) as a meromorphic function with a pole of order $\leq -\kappa_j$. For the case $f = 0$, the solutions u are uniquely determined on the orbits Γ_j by the values of Λ and ϕ at the ends $p_j^-\,,\, p_j^+$ and the number $\alpha_j > 0$. This value is given by the formula

$$u(\Gamma_j) = \epsilon\,\frac{\Lambda(p_j^-)\phi(p_j^+) - \Lambda(p_j^+)\phi(p_j^-)}{i\sin(\pi\alpha_j)} \tag{0.2}$$

with $\epsilon = \pm 1$.

Various properties of the types of vector fields considered here have been studied by many authors and our approach is within the framework of the work contained in [2], [3], [4], [6], [7], [8], [9], [11], [13], [14]. To our knowledge, boundary value problems for elliptic vector fields with interior degeneracies have not been studied, except in [10] where a special case of Problem (0.1) is considered when the boundary functions satisfy $\Lambda(p^-) = \Lambda(p^+)$ and $\phi(p^-) = \phi(p^+)$ on each orbit Γ with ends p^- and p^+. This paper can thus be considered as an outgrowth of Section 12 of [10].

Our basic technique to investigate Problem (0.1) is to make use of the first integrals of L in the connected components of $\Omega\backslash\Gamma_1 \cup \cdots \cup \Gamma_N$ and to reduce the problem to (N+1) classical RH problems for holomorphic functions in the unit disk $\mathbb{D} \subset \mathbb{C}$. However, these new RH problems have discontinuities on $\partial\mathbb{D}$. These types of discontinuous RH problems have been studied by many authors (see in particular the monographs [1], [5], and [12]). In this paper, we adapt the techniques contained in [1].

The organization of this paper is as follows. In Section 1, we recall the main properties of the class of vector fields under consideration. In Section 2, we decompose the open set $\Omega\backslash\Sigma^\infty$ into appropriate components and construct global first integrals for L. The notion of induced indices for a boundary function Λ is introduced in Section 3. In Sections 4 and 5, we study Problem (0.1). Theorem 4.1 gives the general continuous solution of Problem (0.1) when $f = 0$ and $\phi = 0$. Theorem 5.2 deals with the case $f = 0$ and $\phi \neq 0$ and gives the general (meromorphic) solution satisfying (0.2) and in Theorem 5.4, the general case, with $f \neq 0$, is deduced.

1. A CLASS OF VECTOR FIELDS

We summarize some properties of the class of vector fields under consideration. Most of the material listed in this section can be found in [3], [10], and [14].

Let $\widetilde{\Omega}$ be an open subset of $\mathbb{R}^2$ and let L be a vector field given by

$$L = A(x,y)\frac{\partial}{\partial x} + B(x,y)\frac{\partial}{\partial y}\,, \tag{1.1}$$

with A, $B \in C^\omega(\widetilde{\Omega}, \mathbb{C})$, i.e. A and B are real analytic and $\mathbb{C}$-valued in $\widetilde{\Omega}$. We assume that L is nonsingular in $\widetilde{\Omega}$ so that $|A| + |B| > 0$ everywhere in $\widetilde{\Omega}$. We denote by $\overline{L}$ the complex conjugate vector field ($\overline{L} = \overline{A}\partial_x + \overline{B}\partial_y$) and view L as a partial differential operator with symbol $\sigma = A\xi + B\eta$. The operator L is elliptic at a point (x_0, y_0), if $\sigma(x_0, y_0; \xi, \eta) \neq 0$ for every $(\xi, \eta) \in \mathbb{R}^2\backslash 0$. The ellipticity of L at (x_0, y_0) is equivalent to L and $\overline{L}$ being independent at (x_0, y_0). We denote by $\Sigma(\widetilde{\Omega})$ the set of points in $\widetilde{\Omega}$ where L fails to be elliptic. $\Sigma(\widetilde{\Omega})$ is the base projection of the characteristic set of L (we often refer to $\Sigma(\widetilde{\Omega})$ as the characteristic set of L). Consider the $\mathbb{R}$-valued function ϑ defined in $\widetilde{\Omega}$ by

$$\frac{1}{2i}L \wedge \overline{L} = \vartheta(x,y)\frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y} = \mathrm{Im}(A(x,y)\overline{B(x,y)})\frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y}\,. \tag{1.2}$$

Thus, $\Sigma(\widetilde{\Omega})$ is the analytic variety in $\widetilde{\Omega}$ given by

$$\Sigma(\widetilde{\Omega}) = \left\{(x,y) \in \widetilde{\Omega}\,;\;\; \vartheta(x,y) = 0\right\}. \tag{1.3}$$

We assume that L is not a multiple of a real vector field, so that $\Sigma(\widetilde{\Omega})$ is a real analytic variety of dimension ≤ 1. A point $p_0 \in \Sigma(\widetilde{\Omega})$ is of *finite type* if there exists a vector field Y in the Lie algebra generated by L and $\overline{L}$ such that L and Y are independent at p_0. If no such vector field Y exist, the point p_0 is said to be of *inifinite type*. The set of points $p \in \Sigma(\widetilde{\Omega})$ where L is of finite type will be denoted $\Sigma^0(\widetilde{\Omega})$ and those where L is of infinite type will be denoted $\Sigma^\infty(\widetilde{\Omega})$. We have then

$$\Sigma(\widetilde{\Omega}) = \Sigma^\infty(\widetilde{\Omega}) \cup \Sigma^0(\widetilde{\Omega}) \qquad \text{and} \qquad \Sigma^\infty(\widetilde{\Omega}) \cap \Sigma^0(\widetilde{\Omega}) = \emptyset\,.$$

Since the vector field L is nonsingular and real analytic, then it is *locally integrable*. This means that for every point $p \in \widetilde{\Omega}$, there exist an open set $\mathcal{O}_p \subset \widetilde{\Omega}$, $p \in \mathcal{O}_p$, and a function $z_p \in C^\omega(\mathcal{O}_p, \mathbb{C})$ such that $Lz_p = 0$ and $dz_p \neq 0$. The function z_p is a local first integral of L. The use of the local first integrals allows us to obtain local normal forms for the vector field L.

In a neighborhood of an elliptic point $p \in \widetilde{\Omega}\backslash\Sigma(\widetilde{\Omega})$, the vector field L is conjugate to the CR operator. That is, there exist local coordinates (s,t) centered at p in which $z_p = s+it$ is a first integral and L is a multiple of $\dfrac{\partial}{\partial s}+i\dfrac{\partial}{\partial t}$. In a neighborhood of a point of finite type $p_0 \in \Sigma^0(\widetilde{\Omega})$, there exist local coordinates (s,t) centered at p, an integer $k \in \mathbb{Z}^+$, an $\mathbb{R}$-valued and C^ω-function $\phi(s,t)$ satisfying

$$\frac{\partial^j\phi}{\partial t^j}(0) = 0, \;\; j = 1, \cdots k-1, \;\; \frac{\partial^k\phi}{\partial t^k}(0) \neq 0 \tag{1.4}$$

such that $z_p = s + i\phi(s,t)$ is a first integral and L is conjugate to a multiple of

$$\left(1 + i\frac{\partial\phi}{\partial s}\right)\frac{\partial}{\partial t} - i\frac{\partial\phi}{\partial t}\frac{\partial}{\partial s}\,.$$

Moreover, the variety $\Sigma^0(\widetilde{\Omega})$ is locally given by $\{\phi_t = 0\}$. In a neighborhood of a point of infinite type $p \in \Sigma^\infty(\widetilde{\Omega})$, there exist coordinates (s,t) centered at p, an integer $l \in \mathbb{Z}^+$, an $\mathbb{R}$-valued and C^ω-function $\psi(s,t)$ with $\psi(0,t) \not\equiv 0$ such that $z_p = s + is^l\psi(s,t)$ is a first integral and L is conjugate to a multiple of

$$\left(1 + i\frac{\partial s^l\psi}{\partial s}\right)\frac{\partial}{\partial t} - is^l\frac{\partial\psi}{\partial t}\frac{\partial}{\partial s}\,. \tag{1.5}$$

In these local coordinates, $\Sigma^\infty(\widetilde{\Omega})$ is the t-axis ($\{s = 0\}$) and $\Sigma^0(\widetilde{\Omega})$ is given by $\{\psi_t = 0\}$ so that

$$\Sigma(\widetilde{\Omega}) = \{(s,t);\;\; s^l\psi_t(s,t) = 0\;\}\,. \tag{1.6}$$

It follows then that $\Sigma^\infty(\widetilde{\Omega})$ is a one-dimensional real analytic manifold embedded in $\widetilde{\Omega}$ and that L is tangent to $\Sigma^\infty(\widetilde{\Omega})$. For a given open set $U \subset \widetilde{\Omega}$, the connected components of $\Sigma^\infty(\widetilde{\Omega}) \cap U$ will be refereed to as the *orbits* of L in U.

This paper deals with the vector fields with *minimal* orbits (as defined in [10]). That is, we will assume throughout that if $p \in \Sigma^\infty(\widetilde{\Omega})$, then the integer l appearing in the normalization (1.5) is $l = 1$. The semiglobal normalization of vector fields with minimal orbits is given in [10].

A vector field L is said to be *locally solvable* at a point $p \in \widetilde{\Omega}$, if there exists an open set $\mathcal{O} \subset \widetilde{\Omega}$, with $p \in \mathcal{O}$, such that for every function $f \in C_0^\infty(\mathcal{O})$ the equation $Lu = f$ has a distribution solution $u \in \mathcal{D}'(\mathcal{O})$. A vector field L is solvable (and hypoelliptic) at all elliptic points $p \in \widetilde{\Omega}\backslash\Sigma(\widetilde{\Omega})$. The solvability of L at the nonelliptic points $p \in \Sigma(\widetilde{\Omega})$ is given by the Nirenberg-Treves Condition $(\mathcal{P})$. In the case considered here of vector fields in two variables, this condition has a simple formulation. The vector field L is solvable at $p \in \Sigma$ if and only if there exists an open set $\mathcal{O} \subset \widetilde{\Omega}$, with $p \in \mathcal{O}$, such that the function $\vartheta = \mathrm{Im}(A\overline{B})$ defined in (1.2) does not change sign in any connected component of $\mathcal{O}\backslash\Sigma^\infty(\widetilde{\Omega})$. It follows, in particular, that if L is locally solvable at a point of finite type $p \in \Sigma^0(\widetilde{\Omega})$, then the local first integral about p is a local homeomorphism (see [14]). Furthermore, any continuous solution of $Lu = 0$ in a neighborhood of a point $p \notin \Sigma^\infty(\widetilde{\Omega})$ can be written as $u = H \circ z$ where z is a local first integral and where H is a holomorphic function defined in a neighborhood of $z(p) \in \mathbb{C}$. For solutions with isolated singularities, we define a *pole of order s* of a solution u to be a point $p \notin \Sigma^\infty$ such that there exists a meromorphic function M with pole of order s at $z(p)$ such that $u = M \circ z$ in a neighborhood of p. The global solvability of vector fields with minimal orbits is studied in [10].

From now on, we will assume that L satisfies Condition $(\mathcal{P})$ of local solvability and that it has only minimal orbits.

2. Decomposition of Ω and global first integrals

We decompose an open set Ω into suitable components and construct global first integrals for the vector field L. Let L, given by (1.1) with $A, B \in C^\omega(\widetilde{\Omega}, \mathbb{C})$, be such that L satisfies Condition $(\mathcal{P})$ and has only minimal orbits. Let $\Omega \subset \mathbb{R}^2$ be a simply connected and bounded open set with $\overline{\Omega} \subset \widetilde{\Omega}$. For simplicity, we will assume that $\partial\Omega$ is smooth (of class C^∞) and that it is transversal to the manifold $\Sigma^\infty(\widetilde{\Omega})$. Throughout, we will use the following notation

$$\Sigma = \Sigma(\widetilde{\Omega}) \cap \overline{\Omega}, \quad \Sigma^\infty = \Sigma^\infty(\widetilde{\Omega}) \cap \overline{\Omega} \quad \text{and} \quad \Sigma^0 = \Sigma^0(\widetilde{\Omega}) \cap \overline{\Omega}.$$

The set Σ^∞ is therefore a one-dimensional manifold. We denote by Σ_c^∞ the union of the closed curves in Σ^∞ and by Σ_a^∞ the union of the non-closed components of Σ^∞. We have then

$$\Sigma^\infty = \Sigma_c^\infty \cup \Sigma_a^\infty \quad \text{and} \quad \Sigma_c^\infty \cap \Sigma_a^\infty = \emptyset.$$

Hence, Σ_c^∞ consists of finitely many closed curves, each contained in the open set Ω, and Σ_a^∞ consists of finitely many arcs, each connecting two distinct points of $\partial\Omega$ (since $\partial\Omega$ is transversal to $\Sigma^\infty(\widetilde{\Omega})$).

Let $\Gamma_1, \cdots, \Gamma_N$ be the connected components of Σ_a^∞. It is proved in [10] that

$$\Omega\backslash\Sigma_a^\infty = \Omega_1 \cup \cdots \cup \Omega_{N+1}, \tag{2.1}$$

where each component is simply connected and for any two distinct components Ω_j and Ω_k, we have either $\overline{\Omega}_j \cap \overline{\Omega}_k = \emptyset$ or $\overline{\Omega}_j \cap \overline{\Omega}_k = \Gamma_l$, for some component Γ_l of Σ_a^∞. The boundary $\partial\Omega_j$ of each component Ω_j of $\Omega\backslash\Sigma_a^\infty$ is a piecewise smooth curve and consists of an arc in $\partial\Omega$ with ending points on Σ_a^∞, followed by a component of Σ_a^∞, followed by an arc in $\partial\Omega$ and so on. More precisely, if $\Sigma_a^\infty \neq \emptyset$, then for every $j = 1, \cdots, N+1$, there exists an integer $m \in \mathbb{Z}^+$ such that

$$\partial\Omega_j = A_{j1} \cup \Gamma_{j1} \cup \cdots \cup A_{jm} \cup \Gamma_{jm}, \tag{2.2}$$

where the A_{jk}'s are connected components of $\partial\Omega \cap \partial\Omega_j$ and the Γ_{jk}'s connected components of $\partial\Omega_j \cap \Sigma_a^\infty$. Moreover, Γ_{jk} intersects transversally A_{jk} at a point p_{jk}^- and intersects $A_{j(k+1)}$ at a point p_{jk}^+. Hence,

$$\Gamma_{jk} = \operatorname{arc}(p_{jk}^-, p_{jk}^+) \quad \text{and} \quad A_{jk} = \operatorname{arc}(p_{j(k-1)}^+, p_{jk}^-)\,, \tag{2.3}$$

with the understanding that $p_{j0}^+ = p_{jm}^+$.



FIGURE 1. *Decomposition of* Ω *by the* Γ*-orbits*

For each $j \in \{1, \cdots, N+1\}$, let

$$\Omega_j \backslash \Sigma_c^\infty = \Omega_j^0 \cup \widehat{\Omega}_j\,,$$

where Ω_j^0 is the nonrelatively compact component of $\Omega_j \backslash \Sigma_c^\infty$ and $\widehat{\Omega}_j$ is the union of the relatively compact components. Hence, there exist closed orbits $C_1, \cdots, C_p$ of L contained in Ω_j such that

$$\Omega_j = \Omega_j^0 \cup \overline{\operatorname{Int}(C_1)} \cup \cdots \cup \overline{\operatorname{Int}(C_p)}\,, \tag{2.4}$$

where $\operatorname{Int}(C)$ denotes the open domain bounded by the closed curve C. It follows then that

$$\partial\Omega_j^0 = \partial\Omega_j \cup C_1 \cup \cdots \cup C_p\,. \tag{2.5}$$

The following result is proved in [10].

Proposition 2.1. ([10]) *Let* Ω_j *and* Ω_j^0 *be as above with boundaries given by* (2.2) *and* (2.4)*, respectively. Then there exist a function*

$$F_j : \overline{\Omega_j^0} \longrightarrow \overline{\mathbb{D}}\,,$$

where $\mathbb{D}$ *is the unit disk in* $\mathbb{C}$*; points* $z_{j1}, \cdots, z_{jp} \in \mathbb{D}$*; and points* $c_{j1}, \cdots, c_{jm} \in \partial\mathbb{D}$ *such that:*



FIGURE 2. *First integral Z on Ω_2*

- *F_j is C^ω in Ω_j^0 and Hölder continuous on $\overline{\Omega_j^0}$;*
- *$LF_j = 0$ in Ω_j^0 and $dF_j(p) \neq 0$ for every $p \in \Omega_j^0$;*
- *$F_j : \ \Omega_j^0 \longrightarrow \mathbb{D}\backslash\{z_{j1}, \cdots, z_{jp}\}$ is a homeomorphism;*
- *$F_j(C_l) = z_{jl}$ for $l = 1, \cdots, p$ and $F_j(\Gamma_{jk}) = c_{jk}$ for $k = 1, \cdots, m$.*

Remark 2.2. Choose an orientation of $\mathbb{R}^2$ and its induced orientation on the components Ω_j's of $\Omega\backslash\Sigma_a^\infty$. Given a component Ω_j, if the first integral F_j preserves the orientation, then any other first integral also preserves the orientation. The components Ω_j's can thus be divided into two types: Those with orientation preserved and those with orientation reversed by first integrals. Furthermore, if Ω_j and Ω_k are two adjacent components ($\overline{\Omega_j} \cap \overline{\Omega_k} \neq \emptyset$), then one of the component is orientation preserved and the other orientation reversed (see [10] for details).

The first integrals F_j of Proposition 2.1 can be glued together to obtain a global first integral in

$$\Omega^0 = \Omega_1^0 \cup \cdots \cup \Omega_{N+1}^0 \,. \tag{2.6}$$

More precisely, we have the following theorem.

Theorem 2.3. *There exists a function $Z : \overline{\Omega^0} \longrightarrow \overline{\mathbb{D}}$, where Ω^0 is given by* (2.6)*, such that:*

- *$Z \in C^\omega(\Omega^0, \mathbb{D})$, Z is Hölder continuous on $\overline{\Omega^0}$;*

- *$LZ = 0$ and $dZ \neq 0$ in Ω^0;*
- *For each $j = 1, \cdots, N+1$, there exist points $z_{j1}, \cdots, z_{jp} \in \mathbb{D}$ such that*
$$Z(C_l) = z_{jl} \text{ for } l = 1, \cdots, p \quad \text{and} \quad Z: \; \Omega_j^0 \longrightarrow \mathbb{D} \backslash \{z_{j1}, \cdots, z_{jp}\}$$
is a homeomorphism;
- *For each $j = 1, \cdots, N+1$ and $k = 1, \cdots, m$ there exist $c_{jk} \in \partial\mathbb{D}$ such that*
$$Z(\Gamma_{jk}) = c_{jk} \quad \text{and} \quad Z(A_{jk}) = \text{arc}(c_{j(k-1)}, c_{jk}) \, .$$

Proof. Select a component Ω_j of $\Omega \backslash \Sigma_a^\infty$ and label it V_0 so that, as in (2.4), we have

$$V_0 = V_0^0 \cup \overline{\text{Int}(C_1)} \cup \cdots \cup \overline{\text{Int}(C_p)} \, ,$$

where V_0^0 is the non relatively compact component of $V_0 \backslash \Sigma_a^\infty$, and with boundary

$$\partial V_0 = A_1 \cup \Gamma_1 \cup \cdots \cup A_m \cup \Gamma_m \, ,$$

with the A_l's and Γ_l's as in (2.2) and (2.3). Let $F^0 : \overline{V_0^0} \longrightarrow \overline{\mathbb{D}}$ be a first integral of L in V_0^0 satisfying Proposition 2.1. Set, as above, $\Gamma_k = \text{arc}(p_k^-, p_k^+)$, $A_k = \text{arc}(p_{k-1}^+, p_k^-)$, $F^0(C_s) = z_s \in \mathbb{D}$, and $F^0(\Gamma_k) = c_k \in \partial\mathbb{D}$ so that $F^0(A_k) = \text{arc}(c_{k-1}, c_k)$.

For each $k = 1, \cdots, m$ there exists a unique component U_k of $\Omega \backslash \Sigma_a^\infty$ such that $\overline{V^0} \cap \overline{U_k} = \Gamma_k$. Let $F_k : \overline{U_k^0} \longrightarrow \overline{\mathbb{D}}$ be a first integral of L in U_k^0 as in Proposition 2.1. If $F_k(\Gamma_k) = d_k \in \partial\mathbb{D}$, let $\widetilde{F_k} : \overline{U_k^0} \longrightarrow \overline{\mathbb{D}}$ be given by

$$\widetilde{F_k}(x, y) = \frac{c_k}{d_k} F_k(x, y).$$

$\widetilde{F_k}$ is another first integral of L in U_k^0 satisfying Proposition 2.1 and $\widetilde{F_k}(\Gamma_k) = F^0(\Gamma_k) = c_k$. Hence $\widetilde{F_k}$ is a Hölder continuous extension of F^0 to U_k^0 (see [10]). Let

$$V_1 = V_0 \cup U_1 \cup \cdots \cup U_m \, .$$

Define

$$F^1 : \; \overline{V_1^0} = \overline{V_0^0} \cup \overline{U_1^0} \cup \cdots \cup \overline{U_m^0} \longrightarrow \overline{\mathbb{D}}$$

by

$$F^1(x, y) = \begin{cases} F^0(x, y) & \text{if} \quad (x, y) \in \overline{V_0^0}; \\ \widetilde{F_k}(x, y) & \text{if} \quad (x, y) \in \overline{U_k^0} \, . \end{cases}$$

Then, F^1 is a first integral of L in V_1^0 that satisfies the properties of the theorem. This extension of the first integral can be repeated through each orbit contained in ∂V_1 to a larger union of components V_2 and so on until we exhaust all the components of Σ_a^∞ and reach the global first integral in Ω^0. □

3. Induced indices

For a function defined on the boundary of Ω, we define its indices with respect to the components of $\Omega \backslash \Sigma_a^\infty$. Let L and Ω be, respectively, a vector field and an open set as in Section 2. Thus, Ω is simply connected, $\overline{\Omega} \subset \widetilde{\Omega}$, $\partial\Omega$ is a smooth closed curve and transversal to Σ_a^∞. We assume that $\partial\Omega$ is positively oriented (counterclockwise). Let

$$\Omega \backslash \Sigma_a^\infty = \Omega_1 \cup \cdots \cup \Omega_{N+1} \tag{3.1}$$

be the decomposition given in Section 2. We assume that $N \geq 1$ and that Ω_1 is orientation preserved by the first integrals of L. Fix a point $s_0 \in \partial\Omega_1 \backslash \Sigma_a^\infty$ and let $\tau : [0,\ 2\pi] \longrightarrow \partial\Omega$ be a smooth parametrization such that $\tau(0) = \tau(2\pi) = s_0$.

Let $\Lambda : \partial\Omega \longrightarrow \mathbb{T}^1 = \partial\mathbb{D}$ be a Hölder continuous function ($\Lambda \in C^\sigma(\partial\Omega, \mathbb{T}^1)$ with $0 < \sigma < 1$) and denote by $\arg\Lambda$ a continuous branch of the argument of Λ. That is, $\Lambda(s) = \mathrm{e}^{i\arg\Lambda(s)}$ and $\arg\Lambda$ is defined through the parametrization τ as a continuous function

$$\arg\Lambda\circ\tau : [0,\ 2\pi] \longrightarrow \mathbb{R}$$

such that

$$\arg\Lambda(s_0^-) = \arg\Lambda(s_0^+) + \pi q_0 \quad \text{with} \quad q_0 \in \mathbb{Z}\,, \tag{3.2}$$

where $\arg\Lambda(s_0^+) = \lim\limits_{t\to 0^+} \arg\Lambda(\tau(t))$ and $\arg\Lambda(s_0^-) = \lim\limits_{t\to 2\pi^-} \arg\Lambda(\tau(t))$. Note that $q_0/2$ is the winding number of Λ.

Now we define the jump along an orbit $\Gamma \subset \Sigma_a^\infty$ of the function Λ. Such an orbit Γ is the common boundary of two adjacent components Ω_j and Ω_l of $\Omega\backslash\Sigma_a^\infty$. One of the components, say Ω_j is orientation preserved by the first integrals and the other is orientation reversed (Remark 2.2). Consider then Ω_j and its boundary

$$\partial\Omega_j = A_{j1} \cup \Gamma_{j1} \cup \cdots \cup A_{jm} \cup \Gamma_{jm} \tag{3.3}$$

as in (2.2). The orbit appears as $\Gamma = \Gamma_{jk}$ for some k. The boundary $\partial\Omega_j$ inherits its orientation from that of $\partial\Omega$ and so do each of its arcs A_{jl} and Γ_{jl}. Let p_{jk}^- and p_{jk}^+ be the ordered ends of Γ_{jk}. We have

$$\Gamma_{jk} = \mathrm{arc}(p_{jk}^-, p_{jk}^+) \quad \text{and} \quad A_{jk} = \mathrm{arc}(p_{j(k-1)}^+, p_{jk}^-)\,. \tag{3.4}$$

We define the jump of Λ along $\Gamma = \Gamma_{jk}$ as the real number

$$\vartheta_{jk} = \arg\Lambda(p_{jk}^-) - \arg\Lambda(p_{jk}^+)\,. \tag{3.5}$$

To define the index of Λ relative to the component Ω_j, we start by writing the jumps ϑ_{jk} as

$$\vartheta_{jk} = \pi\alpha_{jk} + \pi q_{jk}\,, \tag{3.6}$$

with

$$q_{jk} = \left[\frac{\vartheta_{jk}}{\pi}\right] \in \mathbb{Z} \quad \text{and} \quad \alpha_{jk} = \frac{\vartheta_{jk}}{\pi} - q_{jk} \in [0,\ 1)\,, \tag{3.7}$$

where, for a real number r, $[r]$ denotes the largest integer $\leq r$. Now, we distinguish two types of orbits Γ_{jk}. Those for which the integer q_{jk} is even and those for which it is odd. Let

$$\mathcal{C}_j^1 = \{\Gamma_{jk};\ q_{jk} \in 2\mathbb{Z}+1\}\,, \qquad \mathcal{C}_j^2 = \{\Gamma_{jk};\ q_{jk} \in 2\mathbb{Z}\}\,, \tag{3.8}$$

and let n_j be the number of elements in $\mathcal{C}_j^1$. We define the index κ_j of Λ relative to Ω_j as follows:

$$\begin{aligned} \kappa_j &= \mathrm{Ind}(\Lambda, \Omega_j) = \frac{1}{2}\left(\sum_{k=1}^m q_{jk} - n_j\right) \qquad \text{if } j = 2, \cdots, N+1\,, \\ \kappa_1 &= \mathrm{Ind}(\Lambda, \Omega_1) = \frac{1}{2}\left(q_0 + \sum_{k=1}^m q_{1,k} - n_1\right)\,. \end{aligned} \tag{3.9}$$

Note that $\kappa_j \in \mathbb{Z}$ for $j = 1, \cdots, N+1$.

Example 3.1. Let

$$L=(1+2ixy)\frac{\partial}{\partial y}-i(x^2-1)\frac{\partial}{\partial x}\,. \tag{3.10}$$

It can be seen at once that $L\wedge\overline{L}=-2i(x^2-1)\dfrac{\partial}{\partial x}\wedge\dfrac{\partial}{\partial y}$. Hence, L is elliptic everywhere in $\mathbb{R}^2$ except on the vertical lines $x=1$ and $x=-1$ along which L is tangent. We have then $\Sigma=\Sigma_a^\infty=\{x=\pm1\}$. Furthermore, the orbits $x=1$ and $x=-1$ are minimal. Note that the function $F(x,y)=x+iy(x^2-1)$ is a global first integral of L in $\mathbb{R}^2$.

Let $\Omega=D(0,2)$ be the disk with center 0 and radius 2 in $\mathbb{R}^2$. Then

$$\Omega\backslash\Sigma_a^\infty=\Omega_1\cup\Omega_2\cup\Omega_3\,,$$

where

$$\begin{aligned}\Omega_1\ &=\left\{(x,y)\in\mathbb{R}^2:\ \ 1<x,\ \ x^2+y^2<4\right\}\,;\\ \Omega_2\ &=\left\{(x,y)\in\mathbb{R}^2:\ \ -1<x<1,\ \ x^2+y^2<4\right\}\,;\\ \Omega_3\ &=\left\{(x,y)\in\mathbb{R}^2:\ \ x<-1,\ \ x^2+y^2<4\right\}\,.\end{aligned} \tag{3.11}$$

Σ_a^∞ consists of the vertical segments

$$\Gamma_1=\left\{(1,y):\ -\sqrt{3}\le y\le\sqrt{3}\right\}\quad\text{and}\quad\Gamma_2=\left\{(-1,y):\ -\sqrt{3}\le y\le\sqrt{3}\right\}\,.$$

Since the map $\Phi=(\mathrm{Re}(F),\mathrm{Im}(F))=(x,y(x^2-1))$ is orientation preserving for $|x|>1$ and orientation reversing for $|x|<1$, then Ω_1 and Ω_3 are orientation preserved by first integrals and Ω_2 is orientation reversed. As oriented arcs we have $\Gamma_1=[2\mathrm{e}^{i\pi/3},\ 2\mathrm{e}^{5i\pi/3}]$ and $\Gamma_2=[2\mathrm{e}^{4i\pi/3},\ 2\mathrm{e}^{2i\pi/3}]$.

Let $\Lambda(2\mathrm{e}^{i\theta})=\mathrm{e}^{if(\theta)}$ with $f:\,[0,\ 2\pi]\longrightarrow\mathbb{R}$ smooth and $f(2\pi)=f(0)+\pi q_0$ with $q_0\in2\mathbb{Z}$. The jumps of Λ along Γ_1 and Γ_2 are respectively

$$\vartheta_1=f(\pi/3)-f(5\pi/3)\quad\text{and}\quad\vartheta_2=f(4\pi/3)-f(2\pi/3)\,.$$

The induced indices are respectively

$$\kappa_1=\frac{1}{2}(q_0+q_1-n_1),\quad\kappa_2=\frac{1}{2}(q_1+q_2-n_2)\quad\text{and}\quad\kappa_3=\frac{1}{2}(q_2-n_3)\,,$$

where $q_1=[\vartheta_1/\pi]$; $q_2=[\vartheta_2/\pi]$; $n_1=1$ if q_1 is odd and 0 if not; $n_2=2$ if both q_1 and q_2 are odd, $n_2=1$ if only one q_1 or q_2 is odd and $n_2=0$ if both are even; and $n_3=1$ if q_2 is odd and 0 if not.

In the case $f(\theta)=\theta$, we have $q_0=2$, $\vartheta_1=-4\pi/3$, $\vartheta_2=2\pi/3$, $q_1=-2$, $q_2=0$, $n_1=0$, $n_2=0$ and $n_3=0$. The numbers α_{jk} are in this case $\alpha_1=2/3$ and $\alpha_2=2/3$. The indices are $\kappa_1=0$, $\kappa_2=-1$, and $\kappa_3=0$

In the case $f(\theta)=\pi\sin\theta$, we have $q_0=0$, $\vartheta_1=\pi\sqrt{3}$, $\vartheta_2=-\pi\sqrt{3}$, $q_1=1$, $q_2=-2$, $n_1=1$, $n_2=1$, $n_3=0$, $\alpha_1=\sqrt{3}-1$, $\alpha_2=2-\sqrt{3}$. The indices are $\kappa_1=0$, $\kappa_2=-1$, and $\kappa_3=-1$.

4. The Homogeneous Riemann-Hilbert problem for L

We consider the simplest boundary value problem associated with a vector field. As in the previous section, L is a nonsingular complex vector field with real analytic coefficients given by (1.1) in an open set $\widetilde{\Omega}\subset\mathbb{R}^2$. We assume that L satisfies Condition $(\mathcal{P})$ and that it has only minimal orbits. Let Ω be a bounded, simply connected open set, with $\overline{\Omega} \subset \widetilde{\Omega}$ and with $\partial\Omega$ smooth and transversal to $\Sigma^\infty(\widetilde{\Omega})$. As in the previous sections, we use the decompositions

$$\Omega\backslash\Sigma_a^\infty = \Omega_1 \cup \cdots \cup \Omega_{N+1} \tag{4.1}$$

and for $j = 1, \cdots, N+1$,

$$\partial\Omega_j = A_{j1} \cup \Gamma_{j1} \cup \cdots \cup A_{jm} \cup \Gamma_{jm} \tag{4.2}$$

where Γ_{jk} and A_{jk} are as described in Sections 2 and 3. Let $\Lambda \in C^\sigma(\partial\Omega, \partial\mathbb{D})$ with $0 < \sigma < 1$. For each $j \in \{1, \cdots, N+1\}$, let κ_j be the index of Λ relative to Ω_j and for $k \in \{1, \cdots, m\}$, let $\alpha_{jk} \in [0,\ 1)$ be the associated jump with the orbit Γ_{jk} as defined in (3.7). Denote by δ_j the number of orbits Γ_{jk} with $\alpha_{jk} = 0$. Consider the boundary value problem

$$\begin{cases} Lu = 0 & \text{in } \Omega\,, \\ \operatorname{Re}(\overline{\Lambda} u) = 0 & \text{on } \partial\Omega\,. \end{cases} \tag{4.3}$$

We have the following theorem.

Theorem 4.1. *Problem* (4.3) *has nontrivial continuous solutions on* $\overline{\Omega}$ *that vanish on* Σ_a^∞ *if and only if there exists* $j \in \{1, \cdots, N+1\}$ *with* $2\kappa_j \geq \delta_j$. *Furthermore, the number of independent such solutions is*

$$r = \sum_{\kappa_j \geq \delta_j} (2\kappa_j - \delta_j + 1)\ . \tag{4.4}$$

Proof. For $j = 1, \cdots, N+1$, let $Z_j : \overline{\Omega_j^0} \longrightarrow \overline{\mathbb{D}}$ be the restriction to Ω_j^0 of a first integral Z of L satisfying Theorem 2.3. As in Section 2, Ω_j^0 is the component of $\Omega_j\backslash\Sigma_c^\infty$ given in (2.4) and $\partial\Omega_j^0$ given by (2.5). The boundary $\partial\Omega_j$ is as in (4.2) and $\Gamma_{jk} = \operatorname{arc}(p_{jk}^-, p_{jk}^+)$, $A_{jk} = \operatorname{arc}(p_{j(k-1)}^+, p_{jk}^-)$ (see Section 3), set

$$Z_j(\Gamma_{jk}) = c_{jk} \in \partial\mathbb{D} \quad \text{and} \quad Z_j(A_{jk}) = \operatorname{arc}(c_{j(k-1)}, c_{jk}) \subset \partial\mathbb{D} \tag{4.5}$$

(with $c_{j0} = c_{jm}$). For each $l = 1, \cdots, p$, let $z_l = Z_j(C_l) \in \mathbb{D}$, where C_l is a closed orbit of L contained in $\partial\Omega_j^0$ as in (2.5).

Define the function λ_j on $\partial\mathbb{D}\backslash\{c_{j1}, \cdots, c_{jm}\}$ by $\lambda_j(\zeta) = \Lambda \circ Z_j^{-1}(\zeta)$. Note that λ_j is Hölder continuous on each arc $\overline{Z_j(A_{jk})} \subset \partial\mathbb{D}$. The function $w_j = u \circ Z_j^{-1}$ satisfies the RH-problem

$$\frac{\partial w_j}{\partial \overline{z}} = 0 \quad \text{in } \mathbb{D}\,, \qquad \operatorname{Re}(\overline{\lambda_j}\, w_j) = 0 \quad \text{on } \partial\mathbb{D}\,. \tag{4.6}$$

This is a discontinuous RH-Problem that was considered by several authors (see for instance [1], [5], and [12]). Note that with our definition of the jump ϑ_{jk} of Λ along Γ_{jk}, we have

$$\lambda_j(c_{jk}^-) = \mathrm{e}^{i\vartheta_{jk}} \lambda_j(c_{jk}^+)\,, \tag{4.7}$$

where $\lambda_j(c^\pm) = \lim\limits_{t\to 0^\pm} \lambda_j(c\mathrm{e}^{it})$. Note also that (4.7) is valid whether Z_j is orientation preserving or not.

Now we proceed to the construction of solutions of Problem (4.6) as it is done in Chapter 3 of [1]. We transform the problem into a continuous RH-Problem and use the Schwarz operator to write the general solution. We start by defining the function $\widetilde{\lambda_j}$ on $\partial\mathbb{D}$ by

$$\widetilde{\lambda_j}(\zeta) = \lambda_j(\zeta) \prod_{k=1}^{m} \frac{\overline{(\zeta - c_{jk})}^{\alpha_{jk}}}{|\zeta - c_{jk}|^{\alpha_{jk}}} \tag{4.8}$$

($z^{\alpha_{jk}}$ is defined through any branch of the logarithm). It is verified at once (see [1]) that

$$\widetilde{\lambda_j}(c_{jk}^-) = (-1)^{q_{jk}} \widetilde{\lambda_j}(c_{jk}^+)\,, \tag{4.9}$$

where q_{jk} is the integer associated with the orbit Γ_{jk} and defined through the jump ϑ_{jk} by (3.7). Hence, λ_j is continuous at the points c_{jk} when q_{jk} is even. To obtain a continuous function through the points with q_{jk} odd, we consider the collection $\mathcal{C}_j^1 = \{c_{jk} :\ q_{jk} \text{ odd}\}$ (if not empty) as a set of ordered points in $\partial\mathbb{D}$ (by a the argument function) and write it as

$$\mathcal{C}_j^1 = \{c_{j\mu_s} :\ 1 \le s \le a\}\,, \quad \mu_s < \mu_{s+1} \quad (1 \le s < a \le m)\,. \tag{4.10}$$

We distinguish two cases depending on whether the number of elements, a, in $\mathcal{C}_j^1$ is even or odd. In case a is even, define the arcs in $\partial\mathbb{D}$ with ends on $\mathcal{C}_j^1$ as

$$\begin{aligned} B_s &= \{\zeta \in \partial\mathbb{D} :\ \arg c_{j\mu_s} < \arg\zeta < \arg c_{j\mu_{s+1}}\}\,, \quad 1 \le s < a\,, \\ B_a &= \{\zeta \in \partial\mathbb{D} :\ \arg c_{j\mu_a} < \arg\zeta < \arg c_{j\mu_1} + 2\pi\}\,. \end{aligned} \tag{4.11}$$

Let $\beta : \partial\mathbb{D}\backslash\mathcal{C}_j^1 \longrightarrow \{-1, 1\}$ be the alternating function defined by

$$\beta(\zeta) = (-1)^{s-1} \quad \text{if} \quad \zeta \in B_s \quad 1 \le s \le a\,. \tag{4.12}$$

In the case when a is odd, consider an additional point $c_0 \in \partial\mathbb{D}\backslash\{c_{j1}, \cdots, c_{jm}\}$ and replace $\mathcal{C}_j^1$ by $\widehat{\mathcal{C}_j^1} = \mathcal{C}_j^1 \cup \{c_0\}$ and proceed to define the arcs B_s's and the alternating function β accordingly.

Now we can define the continuous function λ_j^0 on $\partial\mathbb{D}$ by

$$\lambda_j^0(\zeta) = \beta(\zeta)\lambda_j(\zeta) \prod_{k=1}^{m} \frac{\overline{(\zeta - c_{jk})}^{\alpha_{jk}}}{|\zeta - c_{jk}|^{\alpha_{jk}}} \quad \text{if } a \in 2\mathbb{Z}\,, \tag{4.13}$$

$$\lambda_j^0(\zeta) = \beta(\zeta)\lambda_j(\zeta) \frac{\overline{\zeta - c_0}}{|\zeta - c_0|} \prod_{k=1}^{m} \frac{\overline{(\zeta - c_{jk})}^{\alpha_{jk}}}{|\zeta - c_{jk}|^{\alpha_{jk}}} \quad \text{if } a \in 2\mathbb{Z} + 1\,. \tag{4.14}$$

It can be verified ([1]) that such a function λ_j^0 is Hölder continuous on $\partial\mathbb{D}$ and that it has index κ_j.

Assume that a is even, so that the function λ_j^0 is given by (4.13) (the case a odd can be dealt with in a similar fashion). Choose the branch of the logarithm so that $\arg(\zeta - c_{jk}) \to 0$ as $\zeta \to c_{jk}^+$ on $\partial\mathbb{D}$ and $\arg(\zeta - c_{jk}) \to \pi$ as $\zeta \to c_{jk}^-$ on $\partial\mathbb{D}$. Then we have for $k = 1, \cdots, m$

$$\lambda_j^0(c_{jk}) = \beta(c_{jk}^+)\lambda_j(c_{jk}^+) \prod_{l \ne k} \frac{\overline{(c_{jk} - c_{jl})}^{\alpha_{jl}}}{|c_{jk} - c_{jl}|^{\alpha_{jl}}}\,. \tag{4.15}$$

Consider the function w_j^0 defined in $\mathbb{D}$ by

$$w_j^0(z) = \frac{w_j(z)}{\prod_{k=1}^{m}(z - c_{jk})^{\alpha_{jk}}}\,. \tag{4.16}$$

The RH-Problem for w_j^0 is then

$$\frac{\partial w_j^0}{\partial \bar z} = 0 \quad \text{in } \mathbb{D}\,, \qquad \operatorname{Re}(\overline{\lambda_j^0}\, w_j^0) = 0 \quad \text{on } \partial\mathbb{D}\,. \tag{4.17}$$

This is the classical RH-Problem with only the trivial solution, if $\kappa_j < 0$ and with general solution for $\kappa_j \geq 0$, given by

$$w_j^0(z) = z^{\kappa_j} \mathrm{e}^{i\gamma(z)} \left(id_0 + \sum_{l=1}^{\kappa_j} d_l z^l - \overline{d_l} \frac{1}{z^l} \right)\,, \tag{4.18}$$

where $d_0 \in \mathbb{R},\ d_1, \cdots, d_{\kappa_j} \in \mathbb{C}$ are arbitrary constants and where

$$\gamma(z) = \mathcal{S}(\arg(\zeta^{-\kappa_j} \lambda_j^0(\zeta)))(z)\,, \tag{4.19}$$

with $\mathcal{S}$ the Schwarz operator on the disk $\mathbb{D}$:

$$\mathcal{S}(f)(z) = \frac{1}{2\pi i} \int_{\partial\mathbb{D}} f(\zeta)\, \frac{\zeta + z}{\zeta - z}\, \frac{d\zeta}{\zeta}\,. \tag{4.20}$$

Note that for $z = \zeta \in \partial\mathbb{D}$, we have $\operatorname{Re}(\gamma)(\zeta) = \arg \lambda_j^0(\zeta) - \kappa_j \arg \zeta$.

It follows from (4.16) and (4.18) that when $\kappa_j \geq 0$, the general solution

$$w_j(z) = w_j^0(z) \prod_{k=1}^{m} (z - c_{jk})^{\alpha_{jk}} \tag{4.21}$$

of the Problem (4.6), vanishes at each point c_{jk} with $\alpha_{jk} > 0$. At a point $c_{jk_0} \in \partial\mathbb{D}$ with $\alpha_{jk_0} = 0$, the function w_j vanishes only when the coefficients $d_0,\ d_1, \cdots, d_{\kappa_j}$ satisfy the linear equation

$$id_0 + \sum_{l=1}^{\kappa_j} d_l c_{jk_0}^l - \overline{d_l c_{jk_0}^l} = 0\,. \tag{4.22}$$

Since there are δ_j points c_{jk_0} (with $\alpha_{jk_0} = 0$), then the $(2\kappa_j + 1)$-real coefficients $d_0,\ \operatorname{Re}(d_1),\ \operatorname{Im}(d_1),\ \cdots,\ \operatorname{Re}(d_{\kappa_j}),\ \operatorname{Im}(d_{\kappa_j})$ must satisfy δ_j linear equations (over $\mathbb{R}$). We get then $2\kappa_j - \delta_j + 1$ independent solutions if $2\kappa_j \geq \delta_j$ and only the trivial solution, if $2\kappa_j < \delta_j$.

Finally, we can define the continuous solutions of the original Problem (4.3) in Ω by

$$u(x,y) = \begin{cases} 0 & \text{if} \quad (x,y) \in \overline{\Omega_j} \ \text{ with } \ 2\kappa_j < \delta_j\,; \\ w_j(Z_j(x,y)) & \text{if} \quad (x,y) \in \overline{\Omega_j^0} \ \text{ with } \ 2\kappa_j \geq \delta_j\,; \\ w_j(z_l) & \text{if} \quad (x,y) \in \overline{\operatorname{Int}(C_l)} \subset \Omega_j \ \text{ with } \ 2\kappa_j \geq \delta_j\,, \end{cases} \tag{4.23}$$

where w_j is given by (4.21) and the coefficients satisfy (4.22). It follows also that the number of independent solutions is given by (4.4). □

Example 4.2. We continue with the vector L and $\Omega = D(0,2)$ of Example 3.1. We have seen that $\Sigma_a^\infty = \Gamma_1 \cup \Gamma_2$ and $\Omega \backslash \Sigma_a^\infty = \Omega_1 \cup \Omega_2 \cup \Omega_3$.

For $\Lambda(2\mathrm{e}^{i\theta}) = \mathrm{e}^{i\theta}$, we found that $\kappa_1 = \kappa_3 = 0,\ \kappa_2 = -1$, and $\delta_1 = \delta_2 = \delta_3 = 0$. Hence the RH-Problem

$$Lu = 0 \quad \text{in } D(0,2), \qquad \operatorname{Re}\left(\mathrm{e}^{-i\theta} u(2\mathrm{e}^{i\theta})\right) = 0 \quad \text{on } \partial D(0,2)$$

has 2 independent solutions that are identically zero in Ω_2.

For $\Lambda(2\mathrm{e}^{i\theta}) = \mathrm{e}^{i\sin\theta}$, this time we have $\kappa_1 = 0$, $\kappa_2 = \kappa_3 = -1$, and $\delta_1 = \delta_2 = \delta_3 = 0$. Hence the RH-Problem

$$Lu = 0 \quad \text{in } D(0,2), \qquad \mathrm{Re}\left(\mathrm{e}^{-i\sin\theta}u(2\mathrm{e}^{i\theta})\right) = 0 \quad \text{on } \partial D(0,2)$$

has 1 independent solution that is identically zero in $\Omega_2 \cup \Omega_3$.

5. Nonhomogeneous RH problems

In this section we consider nonhomogeneous RH-Problems for the vector field L on a simply connected open set Ω. The assumptions on L and Ω are as in Section 4. The first problem to be studied is

$$\begin{cases} Lu = 0 & \text{in } \Omega\,, \\ \mathrm{Re}\left(\overline{\Lambda}\,u\right) = \phi & \text{on } \partial\Omega\,, \end{cases} \tag{5.1}$$

where the functions Λ and ϕ are Hölder continuous: $\Lambda \in C^\sigma(\partial\Omega, \mathbb{C})$ and $\phi \in C^\sigma(\partial\Omega, \mathbb{R})$, for some $0 < \sigma < 1$. In general such a problem does not have continuous solutions throughout Ω, even in most simple cases, due to the fact that some of the induced indices of Λ might be negative. We will allow then for solutions to have a single pole in each of the regions with negative index (see Section 1 for the definition of a pole). We will also consider only the *generic* case when Λ takes different values at the ends of each arc $\Gamma \in \Sigma_a^\infty$ (so that the corresponding number α defined in (3.7) is positive: $0 < \alpha < 1$). Before, we state the main result about Problem 5.1, we recall the following property of Cauchy type integrals that will be used and whose proof can be found in [12] page 85.

Lemma 5.1. ([12]) *Given a positively oriented curve $\mathcal{A} = \mathrm{arc}(A, B)$ in $\mathbb{C}$; a point $c \in \mathcal{A}$; a function g on $\mathcal{A}$ that is Hölder continuous on $\overline{\mathrm{arc}(A, c)}$ and on $\overline{\mathrm{arc}(c, B)}$; and given a real number $0 < \alpha < 1$, then the Cauchy integral*

$$\Phi(z) = \frac{1}{2\pi i}\int_{\mathcal{A}} \frac{g(\zeta)}{(\zeta - z)^\alpha(\zeta - z)}\, d\zeta \tag{5.2}$$

satisfies the following:

$$\Phi(z) = \frac{e^{i\pi\alpha}g(c^+) - e^{-i\pi\alpha}g(c^-)}{2i\sin(\pi\alpha)}\,\frac{1}{(z-c)^\alpha} + \Phi_0(z)\,, \tag{5.3}$$

for $z \in \mathbb{C}\backslash\mathcal{A}$ to the left of $\mathcal{A}$, with Φ_0 analytic and such that

$$|\Phi_0(z)| \le \frac{K}{|z - c|^{\alpha_0}} \tag{5.4}$$

for some $K > 0$ and $0 < \alpha_0 < \alpha$.

We have the following theorem.

Theorem 5.2. *Let L and Ω be as in* Theorem 4.1 *with $\Omega\backslash\Sigma_a^\infty$ decomposed as in* (2.1) *and $\partial\Omega_j$ given by* (2.2). *Let $\Lambda \in C^\sigma(\partial\Omega, \partial\mathbb{D})$ be generic, $\phi \in C^\sigma(\partial\Omega, \mathbb{R})$ with $0 < \sigma < 1$ and let $\kappa_1, \cdots, \kappa_{N+1}$ be the indices of Λ with respect to $\Omega_1, \cdots, \Omega_{N+1}$, respectively. Then, the Problem* (5.1) *has a solution u such that u is continuous everywhere on $\overline{\Omega}$ except (possibly) at isolated points. The singular points of u consist of a single pole of order $\le -\kappa_j$ in each component Ω_j where $\kappa_j < 0$. Moreover, the value of any such solution u on an orbit $\Gamma = \mathrm{arc}(p^-, p^+)$ is uniquely determined and it is given by*

$$u(\Gamma) = \epsilon\,\frac{\Lambda(p^-)\phi(p^+) - \Lambda(p^+)\phi(p^-)}{i\sin(\pi\alpha)}\,,$$

where $\alpha \in (0,\ 1)$ *is the associated jump of* Λ *along* Γ *and* $\epsilon = 1$ *or* -1*.*

Proof. Let $Z : \overline{\Omega^0} \longrightarrow \overline{\mathbb{D}}$ be the first integral of L as defined in Theorem 2.3. We can assume that $Z^{-1}(0) \cap \Sigma_c^\infty = \emptyset$ (if not replace Z by $H \circ Z$ with H an appropriate automorphism of $\mathbb{D}$). Hence, $Z^{-1}(0)$ consists of $N+1$ points, one point in each component Ω_j^0. For $j \in \{1, \cdots, N+1\}$, let $Z_j : \overline{\Omega_j^0} \longrightarrow \overline{\mathbb{D}}$ be the restriction of Z to $\overline{\Omega_j^0}$ (given by (2.4)) and with $\partial\Omega_j^0$ given by (2.5). As in Theorem 2.3, for $l = 1, \cdots, p$ let $z_{jl} = Z_j(C_l) \in \mathbb{D}$ and for $k = 1, \cdots, m$ let $c_{jk} = Z(\Gamma_{jk}) \in \partial\mathbb{D}$. Let $\lambda_j = \Lambda \circ Z_j^{-1}$ be defined on $\partial\mathbb{D} \backslash \{c_{j1}, \ldots, c_{jm}\}$ and let λ_j^0 be the modification of λ_j as constructed in the proof of Theorem 4.1. Hence, λ_j^0 is Hölder continuous on $\partial\mathbb{D}$ and is defined by (4.13) or (4.14). The function

$$w_j^0(z) = \frac{u \circ Z_j^{-1}(z)}{\prod_{k=1}^m (z - c_{jk})^{\alpha_{jk}}}\,, \qquad z \in \mathbb{D} \tag{5.5}$$

satisfies the nonhomogeneous RH-Problem

$$\frac{\partial w_j^0(z)}{\partial \overline{z}} = 0 \quad z \in \mathbb{D}, \quad \mathrm{Re}\left(\overline{\lambda_j^0}(\zeta) u(\zeta)\right) = \rho(\zeta) \quad \zeta \in \partial\mathbb{D}\,, \tag{5.6}$$

where

$$\rho(\zeta) = \frac{\beta(\zeta)\psi(\zeta)}{\prod_{k=1}^m |\zeta - c_{jk}|^{\alpha_{jk}}}\,, \tag{5.7}$$

with β the alternating function defined in (4.12) and $\psi(\zeta) = \phi \circ Z_j^{-1}(\zeta)$ for $\zeta \in \partial\mathbb{D}$ and $\zeta \neq c_{jk}$.

Suppose first that $\kappa_j \geq 0$. Then, the general solution of Problem (5.6) is given by

$$w_j^0(z) = z^{\kappa_j} \mathrm{e}^{i\gamma(z)} \left(\mathcal{S}(\widehat{\rho})(z) + id_0 + \sum_{l=1}^{\kappa_j} d_l z^l - \overline{d_l}\frac{1}{z^l} \right)\,, \tag{5.8}$$

where $d_0 \in \mathbb{R}$, $d_1, \cdots, d_{\kappa_j} \in \mathbb{C}$ are arbitrary constants, the function $\gamma(z)$ is given by (4.19), $\mathcal{S}$ is the Schwarz operator in $\mathbb{D}$ given by (4.20), and where

$$\widehat{\rho}(\zeta) = \mathrm{e}^{\mathrm{Im}(\gamma(\zeta))}\rho(\zeta)\,. \tag{5.9}$$

We need to understand the behavior of $\mathcal{S}(\widehat{\rho})(z)$ as $z \in \mathbb{D}$ approaches a boundary point c_{jk}. For this we rewrite $\mathcal{S}(\widehat{\rho})(z)$ as

$$\mathcal{S}(\widehat{\rho})(z) = \frac{1}{2\pi i}\int_{\partial\mathbb{D}} \frac{\widehat{\rho}(\zeta)}{\zeta}\,d\zeta + \frac{2z}{2\pi i}\int_{\partial\mathbb{D}} \frac{\widehat{\rho}(\zeta)/\zeta}{\zeta - z}\,d\zeta\,. \tag{5.10}$$

The function $\widehat{\rho}(\zeta)/\zeta$ can be written as

$$\frac{\widehat{\rho}(\zeta)}{\zeta} = \frac{g(\zeta)}{(\zeta - c_{jk})^{\alpha_{jk}}}\,, \tag{5.11}$$

with

$$g(\zeta) = \frac{\mathrm{e}^{\mathrm{Im}(\gamma(\zeta))}\beta(\zeta)\psi(\zeta)}{\zeta \prod_{l \neq k} |\zeta - c_{jl}|^{\alpha_{jl}}} \times \frac{(\zeta - c_{jk})^{\alpha_{jk}}}{|\zeta - c_{jk}|^{\alpha_{jk}}}\,. \tag{5.12}$$

With our choice of a branch of the logarithm as in the proof of Theorem 4.1, we have

$$\begin{aligned} g(c_{jk}^{+}) = &\ \frac{\mathrm{e}^{\mathrm{Im}(\gamma(c_{jk}))}\beta(c_{jk}^{+})\psi(c_{jk}^{+})}{c_{jk}\prod_{l\neq k}|c_{jk}-c_{jl}|^{\alpha_{jl}}}\,; \\ g(c_{jk}^{-}) = &\ \frac{\mathrm{e}^{\mathrm{Im}(\gamma(c_{jk}))}\beta(c_{jk}^{-})\psi(c_{jk}^{-})\mathrm{e}^{i\pi\alpha_{jk}}}{c_{jk}\prod_{l\neq k}|c_{jk}-c_{jl}|^{\alpha_{jl}}}\,. \end{aligned} \tag{5.13}$$

It follows from (5.10), (5.11), (5.12) and Lemma 5.1 that

$$\mathcal{S}(\widehat{\rho})(z) = \frac{\mathrm{e}^{\mathrm{Im}(\gamma(c_{jk}))}}{c_{jk}\prod_{l\neq k}|c_{jk}-c_{jl}|^{\alpha_{jl}}} \times \frac{M_{jk}}{(z-c_{jk})^{\alpha_{jk}}} + \Phi_0(z)\,, \tag{5.14}$$

with

$$|\Phi_0(z)| \leq \frac{K}{|z-c_{jk}|^{\alpha_{jk}^0}} \tag{5.15}$$

for some $K > 0$ and $0 < \alpha_{jk}^0 < \alpha_{jk}$ and where

$$M_{jk} = \frac{\mathrm{e}^{i\pi\alpha_{jk}}\beta(c_{jk}^{+})\psi(c_{jk}^{+}) - \beta(c_{jk}^{-})\psi(c_{jk}^{-})}{2i\sin(\pi\alpha_{jk})}\,. \tag{5.16}$$

Therefore, it follows from (5.8) and (5.14), that the function

$$w_j(z) = w_j^0(z)\prod_{k=1}^{m}(z-c_{jk})^{\alpha_{jk}} \tag{5.17}$$

satisfies (for $z \in \mathbb{D}$ close to c_{jk})

$$w_j(z) = 2z^{\kappa_j+1}\frac{\mathrm{e}^{i\gamma(z)}\mathrm{e}^{\mathrm{Im}(\gamma(c_{jk}))}}{c_{jk}} \times \prod_{l\neq k}\frac{(z-c_{jl})^{\alpha_{jl}}}{|c_{jk}-c_{jl}|^{\alpha_{jl}}} \times M_{jk} + w_j^*(z)\,, \tag{5.18}$$

with w_j^* analytic and $w_j^*(c_{jk}) = 0$. Therefore,

$$w_j(c_{jk}) = \lim_{z\in\mathbb{D}\to c_{jk}} w_j(z) = 2c_{jk}^{\kappa_j}\mathrm{e}^{i\mathrm{Re}(\gamma(c_{jk}))}M_{jk}\prod_{l\neq k}\frac{(c_{jk}-c_{jl})^{\alpha_{jl}}}{|c_{jk}-c_{jl}|^{\alpha_{jl}}}\,. \tag{5.19}$$

Since γ is given by (4.19), then

$$\mathrm{e}^{i\mathrm{Re}(\gamma(c_{jk}))} = c_{jk}^{-\kappa_j}\lambda_j^0(c_{jk}) = c_{jk}^{-\kappa_j}\beta(c_{jk}^{+})\lambda_j(c_{jk}^{+})\prod_{l\neq k}\frac{\overline{(c_{jk}-c_{jl})^{\alpha_{jl}}}}{|c_{jk}-c_{jl}|^{\alpha_{jl}}}\,. \tag{5.20}$$

Hence, using (5.20) and (5.16) in (5.19), we get

$$\begin{aligned} w_j(c_{jk}) &= 2\beta(c_{jk}^{+})\lambda_j(c_{jk}^{+})M_{jk} \\ &= \frac{\lambda_j(c_{jk}^{+})\psi(c_{jk}^{+})\mathrm{e}^{i\pi\alpha_{jk}} - \beta(c_{jk}^{+})\beta(c_{jk}^{-})\lambda_j(c_{jk}^{+})\psi(c_{jk}^{-})}{i\sin(\pi\alpha_{jk})}\,. \end{aligned} \tag{5.21}$$

The expression for $w_j(c_{jk})$ can be further simplified by noticing that

$$\beta(c_{jk}^{+})\beta(c_{jk}^{-}) = (-1)^{q_{jk}} \quad \text{and} \quad \lambda_j(c_{jk}^{-}) = (-1)^{q_{jk}}\mathrm{e}^{i\pi\alpha_{jk}}\lambda_j(c_{jk}^{+}) \tag{5.22}$$

so that

$$w_j(c_{jk}) = (-1)^{q_{jk}}\frac{\lambda_j(c_{jk}^{-})\psi(c_{jk}^{+}) - \lambda_j(c_{jk}^{+})\psi(c_{jk}^{-})}{i\sin(\pi\alpha_{jk})}\,. \tag{5.23}$$

In terms of the values of original functions Λ and ϕ at the ends p_{jk}^- and p_{jk}^+ of the orbit Γ_{jk}, we have

$$w_j(c_{jk}) = (-1)^{q_{jk}} \frac{\Lambda(p_{jk}^-)\phi(p_{jk}^+) - \Lambda(p_{jk}^+)\phi(p_{jk}^-)}{i\sin(\pi\alpha_{jk})}\,. \tag{5.24}$$

In summary, going back to the original problem, we see that in the case $\kappa_j \geq 0$, the function u_j defined in $\overline{\Omega_j}$ by

$$u_j(x,y) = \begin{cases} w_j(Z_j(x,y)) & (x,y) \in \overline{\Omega_j^0}\,, \\ w_j(z_{jl}) & (x,y) \in \mathrm{Int}(C_l), \quad l = 1,\cdots,p\,, \end{cases} \tag{5.25}$$

is continuous on $\overline{\Omega_j}$, satisfies

$$Lu = 0 \ \text{ in } \Omega_j\,, \quad \mathrm{Re}(\overline{\Lambda}(t)u(t)) = \phi(t) \ \ t \in \partial\Omega_j \cap \partial\Omega\,, \tag{5.26}$$

and is constant on each arc Γ_{jk} along which it is uniquely determined by $u(\Gamma_{jk}) = w_j(c_{jk})$ given by (5.24).

Now we consider the case $\kappa_j < 0$. The solution

$$w_j^0(z) = z^{\kappa_j} \mathrm{e}^{i\gamma(z)} \mathcal{S}(\widehat{\rho})(z) \tag{5.27}$$

of Problem (5.6) is meromorphic with a single pole at $z = 0$ of order $\leq -\kappa_j$. The function defined in $\overline{\Omega_j} \backslash Z_j^{-1}(0)$ by

$$u_j(x,y) = \begin{cases} w_j(Z_j(x,y)) & (x,y) \in \overline{\Omega_j^0} \backslash Z_j^{-1}(0)\,, \\ w_j(z_{jl}) & (x,y) \in \mathrm{Int}(C_l), \quad l = 1,\cdots,p\,, \end{cases} \tag{5.28}$$

is continuous with a pole of order $\leq -\kappa_j$ at the point $Z_j^{-1}(0)$ and solves problem (5.26) in $\Omega_j \backslash Z_j^{-1}(0)$. Furthermore, its value on each orbit Γ_{jk} is again given by (5.24).

These various functions u_j defined in Ω_j yield a well defined continuous function u on the whole open set Ω (except at isolated poles). Furthermore, it is easily verifiable that u solves the RH-Problem (5.1). □

Remark 5.3. The RH-Problem (5.1) could admit continuous solutions throughout Ω even when some of the indices $\kappa_j < 0$, provided that the boundary functions Λ and ϕ satisfy $(-\kappa_j - 1)$-conditions on $\partial\Omega_j$. These conditions simply mean that $\mathcal{S}(\widehat{\rho})$ has a zero of order $\geq -\kappa_j$ at $z = 0$ (where $\widehat{\rho}$ is given by (5.9)). That is, $\displaystyle\int_{\partial\mathbb{D}} \widehat{\rho}(\zeta)\zeta^{-s} d\zeta = 0$ for $s = 1,\cdots,-\kappa_j$. In terms of integrals over $\partial\Omega_j$, we get the conditions

$$\sum_{k=1}^{m} \int_{A_{jk}} \widehat{\rho}(Z_j(t)) \frac{dZ_j(t)}{Z_j^s(t)} = 0 \quad s = 1,\cdots,-\kappa_j\,. \tag{5.29}$$

The last problem we consider in this paper is

$$\begin{cases} Lu = f & \text{in } \ \Omega\,, \\ \mathrm{Re}\left(\overline{\Lambda}\, u\right) = \phi & \text{on } \ \partial\Omega\,. \end{cases} \tag{5.30}$$

We assume that Λ and ϕ are as in Theorem 5.2 and $f \in C^\infty(\overline{\Omega})$ has zero periods on the closed orbits of L. By this, we mean that if we consider the dual differential form $\omega = Bdx - Ady$ of the vector field L, then there exists a C^∞-differential form

η in $\overline{\Omega}$ such that $f dx\wedge dy=\omega\wedge\eta$, and we say that f has *zero periods* on the closed orbits of L if

$$\int_C \eta = 0 \quad \text{for every closed orbit} \quad C\subset\Sigma_c^\infty\,. \tag{5.31}$$

It should be noted that this condition depends only on the function f and not on the particular choice of its representative η (any other representative is of the form $\eta+g\omega$ for some function g). It is proved in [10] that condition (5.31) is a necessary and sufficient condition for the global solvability of L on Ω.

Now we can state our result regarding Problem (5.30).

Theorem 5.4. *Let* $L\,,\ \Omega\,,\ \Lambda\,,\ \phi$ *be as in* Theorem 5.2. *Let* κ_j *be the index of* Λ *relative to the component* Ω_j *of* $\Omega\backslash\Sigma_a^\infty$ *and* $f\in C^\infty(\overline{\Omega})$ *satisfies condition* (5.31). *Let* $\kappa_{j_1},\cdots,\kappa_{j_n}$ *be the negative indices of* Λ. *Then, for every* $l=1,\cdots,n$ *there exists a point* $s_{j_l}\in\Omega_{j_l}\backslash\Sigma^\infty$ *such that Problem* (5.30) *has a solution* u *that is Hölder continuous on* $\overline{\Omega}\backslash\{s_{j_1},\cdots,s_{j_n}\}$ *and* C^∞ *in* $\overline{\Omega}\backslash\Sigma^\infty\cup\{s_{j_1},\cdots,s_{j_n}\}$. *Furthermore, for each* s_{j_l} *there exist an open set* $\mathcal{O}_{j_l}$, *with* $s_{j_l}\in\mathcal{O}_{j_l}\subset\Omega$, *and a function* $m_{j_l}\in C^\infty(\mathcal{O}_{j_l})$ *such that* $u-m_{j_l}$ *has a pole of order* $\le-\kappa_{j_l}$ *at* s_{j_l}.

Proof. We know from [10] that for a given $f\in C^\infty(\overline{\Omega})$ satisfying (5.31), there exists a function v, Hölder continuous on $\overline{\Omega}$ and in $C^\infty(\overline{\Omega}\backslash\Sigma_a^\infty)$ such that $Lv=f$. Theorem 5.2 implies that the RH-Problem

$$Lw=0 \ \text{ in } \Omega\,, \qquad \mathrm{Re}(\overline{\Lambda}\,w)=\phi-\mathrm{Re}(\overline{\Lambda}\,v)$$

has a solution w such that w is Hölder continuous on $\overline{\Omega}\backslash\{s_{j_1},\ldots,s_{j_n}\}$, is C^ω in $\overline{\Omega}\backslash\Sigma^\infty\cup\{s_{j_1},\ldots,s_{j_n}\}$, and has a pole of order $\le-\kappa_{j_l}$ at the points s_{j_l}. The function $u=w+v$ is then the sought solution of Problem (5.30). □

References

1. H. Begehr, *Complex analytic methods for partial differential equations. An introductory text*, World Scientific Publishing Co., Inc., NJ (1994). MR1314196(95m:35031)
2. A. Bergamasco and A. Meziani, *Solvability near the characteristic set for a class of planar vector fields of infinite type*, Ann. Inst. Fourier, **55**, (1) (2005), 77-112. MR2141289(2005m:35028)
3. S. Berhanu, P. Cordaro, and J. Hounie, *An introduction to involutive structures*, New Math. Mono., **6**, Cambridge University Press, Cambridge, (2008). MR2397326(2009b:32048)
4. S. Berhanu and A. Meziani, *Global properties of a class of planar vector fields of infinite type*, Comm. Partial Diff. Equ. **22** (1-2) (1997),99-142. MR1434140(98c:35003)
5. F. Gakhov, *Boundary Value Problems*, Dover Publ. Inc., New York (1966). MR1106850(92b:45005)
6. J. Hounie, *Globally hypoelliptic and globally solvable first-order evolution equations*, Trans. Amer. Math. Soc., **252** (1979), 233-248. MR0534120(80h:35035)
7. J. Hounie, *A note on global solvability of vector fields*, Proc. Amer. Math. Soc., **94**, 1 (1985), 61-64. MR0781057(86h:35032)
8. A. Meziani, *On planar elliptic structures with infinite type degenerecy*, J. Funct. Analysis, **179** (2001), 333-373. MR1809114(200k:35122)
9. A. Meziani, *Elliptic planar vector fields with degeneracies*, Trans. Amer. Math. Soc., **357** (2005), 4225-4248. MR2159708(2006f:35107)
10. A. Meziani, *Global solvability of real analytic complex vector fields in two variables*, J. Diffirential Equations, **251** (2011), 2896-2931. MR2831718(2012g:35031)
11. A. Meziani, *On first and second order planar elliptic equations with degeneracies*, Memoirs of the Amer. Math. Soc., Vol. 217, **1019** (2012)
12. N. Muskhelishvili, *Singular Integral Equations*, Dover Pbubl. Inc., New York (1992). MR1215485(94a:45001)

13. L. Nirenberg and F. Treves, *Solvability of a first-order linear differential equation*, Comm. Pure Appl. Math., **16** (1963), 331-351. MR0163045(29#348)
14. F. Treves, *Hypo-Analytic Structures: Local Theory*, Princeton Mathematical Series, **40**, Princeton Univ. Press, NJ (1992). MR1200459(94e:35014)

[1]Lab. AMNEDP, Department of Mathematics, USTHB, Po Box 32 El Alia, Bab Ezzouar 16111 Algiers, Algeria

[2]Department of Mathematics, Florida International University, 11200 S.W. 8th Street, Miami, FL, 33199, USA e-mail: meziani@fiu.edu